

Kumulus[®] WG

Reg. No. L2404; N-AR 0732

Act No. 36 of 1947

**READ ATTACHED PACKAGE OR LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

GROUP	M02	FUNGICIDE
-------	-----	-----------

A water dispersible granular fungicide and acaricide, for the control of powdery mildew, various other fungal diseases, as well as certain mites and thrips on various crops, as listed. Also for the control of pernicious scale, on dormant roses.

‘n Waterdispergeerbare korrelswamdoder en -mytdoder, vir die beheer van poeieragtige meeldou, verskeie ander swamsiektes, asook sekere myte en blaaspootjies op verskeie gewasse, soos aangedui. Ook vir die beheer van verderflike dopluis, op rustende rose.

If medical advice is needed, have product container or label at hand.

Active ingredients / Aktiewe bestanddele:

Sulphur.....800 g/kg

Registered by:

BASF South Africa (Pty) Ltd
Co. Reg. No. 66/10235/07
852 Sixteenth Road
P.O. Box 2801
Halfway House 1685
+27 11 203 2400

Geregistreeer deur:

BASF Suid-Afrika (Edms) Bpk
Mpy Reg. Nr 66/10235/07
852 Sestiende Laan
Posbus 2801
Halfweghuis 1685
+27 11 203 2400

Batch No. & Date manufactured: Refer to details printed on container
Lotnommer & Datum vervaardig: Verwys na besonderhede gedruk op houer

EMERGENCY NUMBER / NOODNOMMER: +27 82 446 8946

25 kg

ENGLISH

WARNINGS

ALLOW THE MINIMUM NUMBER OF WEEKS BETWEEN LAST APPLICATION AND HARVEST:

Wine grapes	4 – 6
Table grapes	2
All other crops	no waiting period required

- Store in a cool, dry place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.
- Flammable – do not store near open flames or sparks.
- RE-ENTRY INTERVAL: Do not enter treated area until spray deposit has dried unless wearing protective clothing

Although this remedy has been tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors, such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore, does not accept responsibility for damage to crops, vegetation, the environment, harm to man or animal, or for lack of performance of the remedy concerned, due to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

Precautionary Statements

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read label before use.
- Do not inhale spray mist.
- Destroy empty bags and never use for any other purpose.
- Prevent contamination of food, feed, drinking water and eating utensils.
- **TRIPLE RINSE THE EMPTY CONTAINER AS FOLLOWS:** Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za). Do not bury, burn, or donate the container to any other parties that may use it as a container for food or beverages.

RELEVANT SUBSTANCES

Hazardous ingredients According to UN GHS criteria	CAS number	Concentration (W/W)
Sulphur	7704-34-9	80 %
Lignosulfonic acid, sodium salt	8061-51-6	< 20 %

FIRST AID TREATMENT

Remove contaminated clothing.

ON CONTACT WITH EYES:	Wash affected eyes for at least 15 minutes under running water with eyelids held open.
ON INGESTION:	Rinse mouth and then drink plenty of water.
ON SKIN CONTACT:	Wash thoroughly with soap and water.
IF INHALED:	Keep patient calm, remove to fresh air.

Most important symptoms and effects, both acute and delayed

Symptoms: No significant reaction of the human body to the product known.

Indication of any immediate medical attention and special treatment needed

Treatment: Symptomatic treatment (decontamination, vital functions).

RESISTANCE WARNING

Generally considered as a low risk group, without signs of resistance developing fungicides. No cross resistance occur between group members M1 – M9. Kumulus® WG is a FRAC Group Code M2 fungicide, for resistance management. Any fungal population may contain individuals naturally resistant to Kumulus® WG and other Group Codes M1 – M9 fungicides. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly and exclusively in programmes. These resistant fungi may not be controlled by Kumulus® WG or any other Group Codes M1 – M9 fungicides.

To delay fungicide resistance:

- **Avoid exclusive repeated use of fungicides from the same Fungicide Group Code. Alternate or tank-mix with products from different Fungicide Group Codes.**
- **For tank mixing or alternation with products in other Fungicide Group Codes, refer to applicable individual product labels.**
- **Integrate other control methods (chemical, cultural, biological), into disease control programmes.**

FOR SPECIFIC INFORMATION ON RESISTANCE MANAGEMENT, CONTACT THE REGISTRATION HOLDER, BASF SA (Pty) Ltd.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	PEST/ DISEASE	DOSAGE g/100 ℓ	DIRECTIONS FOR APPLICATIONS
AVOCADO	Thrips (Thripidae spp.)	300	Apply when first symptoms are noticed.
PAPAYA	Powdery mildew (<i>Ovulariopsis papayae</i>)	300	Apply when first symptoms are noticed. Repeat weekly.
CITRUS			Full cover spray. Allow four weeks safety period between sulphur- and oil treatments.
	Flat mite (<i>Brevipalpus californicus</i>)	150	September – May
	Grey mite (<i>Calcarus citrifolii</i>)	200	During winter or at blossom drop.
	Silver mite (<i>Hemitarsonemus latrus</i>)	250	Apply in spring if russeting occurred in previous season.
	Rust mite (<i>Phyllocoptruta oleivora</i>)	300	December – May. Spray whole orchard.
	Thrips (<i>Scirtothrips aurantii</i>)	300	Apply at 80 – 100% petal drop. Repeat after 10 days if necessary.
APPLES AND PEARS	Budmite (<i>Eriophyes pyri</i>)	400	High volume full cover spray at closed bud stage.
	Powdery mildew (<i>Podosphaera leucotricha</i>) and Scab (<i>Venturia inaequalis</i>)	200 – 400	Pre-blossom stage: 300 – 400 g/100 ℓ Early post-blossom stage: 250 g/100 ℓ Late post-blossom stage: 200 g/100 ℓ Do not apply on sulphur sensitive cultivars.
PEACHES, PLUMS, APRICOTS	Rust (<i>Tranzschelia</i> spp.)	150 – 300	Winter Rainfall region: Start applications early November. Summer Rainfall Region: Apply from middle February, or as soon as first symptoms are noticed.
PEACHES	Powdery mildew (<i>Oidium leucoconium</i>)	200 – 300	Apply at first signs of infection. Repeat every 10 – 14 days if weather favours disease. Dosage depends on severity of disease.

GRAPE VINES	Powdery mildew (<i>Uncinula necator</i>)	300 not less than 3 kg/ha	Winter Rainfall Region: Start application when shoots are 2,5 cm long. Summer Rainfall Region: Start application mid-summer, particularly during drought, or when first symptoms are noticed. Warning: Heavy applications may cause scorching, especially on warm days. Do not apply within 14 days before harvest of table grapes and 4 – 6 weeks in case of wine grapes (pea size). 4 – 6 weeks in case of wine grapes (pea size). Use in a minimum of 500 ℓ water /ha. Do not apply when temperature is above 32° C
ORNAMENTALS	Thrips (Thripidae spp.) (Thripidae spp.)	400	Full cover spray. Sulphur may injure some ornamentals.
	Red Spider Mite [<i>Tetranychus urticae</i> (<i>cinnabarinus</i>)] Powdery mildew (<i>Oidium</i> spp.)	300	
BEANS	Thrips (<i>Scirtothrips occipitalis</i>) Red Spider Mite [<i>Tetranychus urticae</i> (<i>cinnabarinus</i>)]	400	Full cover spray. Repeat at 10 – 14 day intervals.
TOMATOES	Rust mite (<i>Aculus lycopersicae</i>)	400	Full cover spray. Repeat at 2 – 4 weeks intervals.
CUCURBITS	Powdery mildew (<i>Oidium erysiphoides</i>)	300	Do not use on muskmelon and cucumbers. Start application at first signs of disease and repeat every 7 – 14 days. Depending on foliage apply up to 1500 ℓ spray mixture / ha. Ensure total coverage of all parts of the plant.
PEAS	Powdery mildew (<i>Erysiphe pisi</i>)	300	Apply when first symptoms of disease appear and repeat every 7 – 10 days. Apply up to 1000 ℓ spray mixture / ha and ensure good coverage.
ROSES Flower Carpet Blossom Magic Iceberg Sharifa Double Delight	Pernicious scale (<i>Quadraspidiotus perniciosus</i>)	2400 (120 g/5 l)	Full cover spray (Apply to dormant plants only, that is plants without leaves). Also for home garden use.

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of this product in a manner inconsistent with its labeling, all of which are beyond the control of BASF South Africa (Pty) Ltd. ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of the Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

AFRIKAANS

WAARSKUWINGS

LAAT DIE VOLGENDE MINIMUM AANTAL WEKE TOE TUSSEN LAASTE TOEDIENING EN OES

Wyndruiwe	4 – 6
Tafeldruiwe	2
Alle ander gewasse	geen wagperiode word vereis nie.

- Stoor in 'n koel, droë plek weg van kos en voer.
- Hou buite bereik van kinders, oningelgte persone en diere.
- Brandbaar – moenie naby oop vuur of vonke stoor nie.
- HERBETREDING: Moet nie behandelde gebied betree alvorens die spuitneerslag droog is nie, tensy beskermende oorklere gedra word.

Alhoewel hierdie middel onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan, deur faktore soos abnormale grond-, klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie, en die voorkoms van weerstand van die siekte teen die betrokke middel beïnvloed kan word, die metode, tyd en akkuraatheid van toediening kan ook die werking en doeltreffendheid van die middel beïnvloed. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei en die omgewing nie, of vir nadelige effek op mens of dier nie, of vir onvoldoende prestasie van die betrokke middel weens die versuim van die gebruiker om etiketaanwysings na te kom nie, of weens toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in geval van enige onsekerheid.

VOORSORGMATREËLS

Voorsorgverklarings

- Indien mediese advies benodig word, hou die produkhouer of -etiket byderhand.
- Hou buite bereik van kinders.
- Lees sorgvuldig en volg alle instruksies.
- Moenie spuitnewel inasem nie.
- Vernietig leë sakke en moet nooit vir enige ander doel gebruik nie.
- Vermo besoedeling van voedsel, voer, eetgerei en drinkwater.
- SPOEL LEË HOUER DRIE KEER AS VOLG: Keer die leë houer oor die spuit of mengtenk om en dreineer vir minstens 30 sekondes nadat die vloei tot 'n gedrup verminder het. Spoel daarna die leë houer drie keer na mekaar met een kwart van die houervolume vars water en giet die spoelwater in die spuittenk of mengtenk oor Kap gate in die drie keer gespoelde houer waarna dit aan 'n goedgekeurde versamelaar of verwerker oorhandig moet word (www.croplife.co.za). Moet nie die houer begrawe, verbrand of aan enige ander partye skenk wat dit as houer vir voedsel of drinkgoed mag gebruik nie.

RELEVANTE STOWWE

Gevaarlike Bestanddele Volgens UN GHS kriteria	CAS number	Concentration (W/W)
Sulphur	7704-34-9	80 %
Lignosulfonic acid, sodium salt	8061-51-6	< 20 %

NOODHULPBEHANDELING

Verwyder gekontamineerde klere

INDIEN IN DIE OË:	Spoel aangetaste oë vir ten minste 15 minute met lopende water, terwyl die ooglede oopgehou word. .
INDIEN INGESLUK:	Spoel mond dadelik uit en drink baie water.
INDIEN OP DIE VEL:	Was aangetaste area deeglik met seep en water.
INDIEN INGEASEM:	Hou pasiënt kalm en verwyder na vars lug.

Indikasie van enige onmiddellike mediese aandag en spesiale behandeling nodig

Behandeling: Behandel simptome (reinig, belangrikste funksies). Daar is geen spesifieke teenmiddel nie.

WAARSKUWING TEEN WEERSTAND

Swamdoders M1 – M9, word beskou as 'n lae risiko groep, met geen tekens van weerstand wat ontwikkel teen ander swamdoders nie. Geen kruisweerstand is aangemeld tussen die swamdoders in hierdie groepe nie. Kumulus® is 'n FRAC Groepkode M2 swamdoder. Enige populasie van 'n spesifieke swam mag individue insluit wat 'n natuurlike weerstand teen Kumulus®, of enige ander Groepkodes M1 – M9 swamdoders het. Indien hierdie swamdoders in herhaaldelike en eksklusiewe programme aangewend word, kan die weerstandbiedende individue uiteindelik die swampopulasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur Kumulus® of enige ander Groepkodes M1 – M9 swamdoders, beheer word nie.

Om weerstand teen swamdoders te vertraag:

- Vermo die eksklusiewe herhaaldelike gebruik van swamdoders in dieselfde Swamdodergroepkode. Wissel af met, of gebruik tenkmengels van produkte in verskillende Swamdodergroepkodes.
- Sien toepaslike individuele produketikette wanneer daar afgewissel word, of tenkmengels gebruik word, met produkte in ander Swamdodergroepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies), in swamdoderprogramme.

VIR SPESIFIEKE INLIGTING OOR WEERSTANDBESTUUR, KONTAK DIE REGISTRASIEHOUER, BASF SA (Edms) Bpk.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS VOORGESKRYF

GEWAS	PLAAG / SIEKTE	DOSIS g/100 ℓ g/100 ℓ	GEBRUIKSAANBEVELINGS
AVOKADO	Blaaspootjies (<i>Thripidae</i> spp.)	300	Dien toe sodra eerste simptome verskyn.
PAPAJA	Poeieragtige meeldou (<i>Ovulariopsis papayae</i>)	300	Dien toe sodra eerste simptome verskyn. Herhaal weekliks.
SITRUS			Volle dekteoediening. Laat vier weke veiligheidsperiode verstryk tussen swael- en oliebehandelings.
	Platmyt (<i>Brevipalpus californicus</i>)	150	September – Mei.
	Grysmyt (<i>Calcarus citrifolii</i>)	200	In die winter of by blomblaarval.
	Silwermyt (<i>Hemitarsonemus latrus</i>)	250	Dien in lente toe as silwer-verruwing die vorige seisoen voorgekom het.
	Roesmyt (<i>Phyllocoptruta oleivora</i>)	300	Desember – Mei. Spuit die hele boord.
	Blaaspootjie (<i>Scirtothrips aurantii</i>)	300	Dien by 80 – 100% blomblaarval toe. Herhaal na 10 dae indien nodig.
APPELS EN PERE	Knopmyt (<i>Eriophyes pyri</i>)	400	Hoë volume volle dekteoediening in toeknopstadium.
	Poeieragtige meeldou (<i>Podosphaera leucotricha</i>) en Skurfsiekte (<i>Venturia inaequalis</i>)	200 – 400	Voor bloeistadium: 300 – 400 g/100 ℓ Vroeë nabloeistadium: 250 g/100 ℓ Laat nabloeistadium: 200 g/100 ℓ Moenie op swael-sensitiewe kultivars spuit nie.
PERSKES, PRUIME, APPELKOSE	Bruinroes (<i>Tranzhelia</i> spp.)	150 – 300	Winterreëvalstreek: Begin toedien vroeg November. Somerreëvalstreek: Dien vanaf middel Februarie toe, of sodra eerste simptome opgemerk word.
PERSKES	Poeieragtige meeldou (<i>Oidium leucoconium</i>)	200 – 300	Dien toe sodra eerste simptome verskyn. Herhaal met 10 – 14 dae tussenposes, solank die weer die siekte begunstig. Dosis hang af van omvang van besmetting.

WINGERD	Poeieragtige meeldou (<i>Uncinula necator</i>)	300 nie minder as 3 kg/ha nie	Winterreëvalstreek: Begin toedien sodra lote 2,5 cm lank is. Somerreëvalstreek: Begin toedien middel somer veral as dit droog is, of sodra eerste simptome voorkom. Waarskuwing: Swaar toediening van swael mag brandskade veroorsaak, veral op warm dae. Moenie tafeldruive binne 14 dae voor oes, en wyndruive binne 4 – 6 weke (ertjegrootte) voor oes behandel nie. Moenie in minder as 500 ℓ water /ha spuit nie. Moenie spuit wanneer temperatuur hoër as 320° C is nie.
BLOMME EN SIERPLANTE	Blaaspootjies (<i>Thripidae</i> spp.)	400	Volle dekteoediening. Swael mag sekere sierplante brand.
	Rooispinmyt [<i>Tetranychus urticae</i> (<i>cinnabarinus</i>)] Poeieragtige meeldou (<i>Oidium</i> spp.)	300	
BONE	Blaaspootjies (<i>Scirtothrips occipitalis</i>) Rooispinmyt [<i>Tetranychus urticae</i> (<i>cinnabarinus</i>)]	400	Volle dekteoediening. Herhaal elke 10 – 14 dae
TAMATIES	Roesmyt (<i>Aculus lycopersicae</i>)	400	Volle dekteoediening. Herhaal elke 2 – 4 weke.
PAMPOEN-GEWASSE	Poeieragtige meeldou (<i>Oidium erysiphoides</i>)	300	Moenie op komkommers en spanspekke gebruik nie. Begin toedien sodra eerste simptome verskyn. Herhaal met 7 – 14 dae tussenposes. Dien tot 1500 ℓ spuitmengsel / ha toe afhangende van blaardigtheid. Verseker deeglike benatting van alle plantdele.
ERTJIES	Poeieragtige meeldou (<i>Erysiphe pis</i>)	300	Dien toe wanneer eerste simptome verskyn en herhaal elke 7 – 10 dae. Dien tot 1000 ℓ spuitmengsel / ha toe en verseker deeglike benatting.
ROSE Flower Carpet, Blossom Magic, Ice Berg, Sharifa, Double Delight	Verdeflike dopluis (<i>Quadraspidiotus perniciosus</i>)	2400 (120 g/5 l)	Volle dekteoediening. (Dien slegs op rustende plante toe, dus plante sonder blare). Ook vir huistuin gebruik.

VERKOOPSVORWAARDES EN WAARBORG

Die GEBRUIKSAANWYSINGS van hierdie produk weerspieël die opinie van kundiges, gebaseer op gebruik in die veld en in proewe. Die aanwysings word as betroubaar geag en moet noukeurig nagekom word. Dit is egter onmoontlik om alle risiko's wat inherent met die gebruik van hierdie produk geassosieer word, uit te skakel. Gewasskade, oneffektiwiteit of ander onopsetlike gevolge mag voorkom as gevolg van faktore soos weerstoestand, teenwoordigheid van ander materiale, of wanneer die produk op 'n wyse wat teenstrydig met die etiket is, gebruik word; dit alles is buite die beheer van BASF Suid-Afrika (Edms) Bpk ("BASF") of die Verkoper. In die mate waarin dit in ooreenstemming met toepaslike wetgewing is, sal alle risiko's deur die Koper aanvaar word.

BASF waarborg dat hierdie produk aan die vereistes van die chemiese beskrywing op die etiket voldoen, en is redelikerwys geskik vir die gebruiksdoeleindes soos uiteengesit in die GEBRUIKSAANWYSINGS, onderhewig aan die inherente risiko's, soos hierbo na verwys.

IN DIE MATE WAT DIT MET TOEPASLIKE WETGEWING OOREENSTEM, GEE BASF GEEN ANDER UITDRUKKING OF GEÏMPLISEERDE WAARBORG VAN GESKIKTHEID OF VERKOOPSVERNUF OF ENIGE ANDER UITDRUKKING OF GEÏMPLISEERDE WAARBORG NIE.

IN DIE MATE WAT DIT MET TOEPASLIKE WETGEWING OOREENSTEM, SAL DIE KOPER SE UITSLUITLIKE REGSMIDDEL EN BASF SE UITSLUITLIKE AANSPREEKLIKHEID, HETSY IN KONTRAK, ONREGMATIGHEID, NALATIGHEID, STRENG AANSPREEKLIKHEID, OF ANDERSINDS, BEPERK WORD TOT TERUGBETALING VAN DIE AANKOOPPRYS VAN DIE PRODUK.

IN DIE MATE WAT DIT MET TOEPASLIKE WETGEWING OOREENSTEM, DOEN BASF EN DIE VERKOPER, AFSTAND VAN ENIGE AANSPREEKLIKHEID TEN OPSIGTE VAN VOORTSPRUITENDE SKADE, SKADEVERGOEDING, SPESIALE OF INDIREKTE SKADE, UIT DIE GEBRUIK OF HANTERING VAN HIERDIE PRODUK.

BASF en die Verkoper bied hierdie produk aan, en die Aankoper en Gebruiker aanvaar dit, onderhewig aan die voorafgaande Verkoopsvoorwaardes en Waarborg, wat slegs gewysig mag word deur 'n skriftelike ooreenkoms, wat na behore deur 'n gemagtigde verteenwoordiger van BASF onderteken is.

Kumulus[®] WG

Reg. No. L2404; N-AR 0732

Act No. 36 of 1947

**READ ATTACHED PACKAGE OR LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

GROUP	M02	FUNGICIDE
-------	-----	-----------

A water dispersible granular fungicide and acaricide, for the control of powdery mildew, various other fungal diseases, as well as certain mites and thrips on various crops, as listed. Also for the control of pernicious scale, on dormant roses.

'n Waterdispergeerbare korrelswamdoder en -mytdoder, vir die beheer van poeieragtige meeldou, verskeie ander swamsiektes, asook sekere myte en blaaspootjies op verskeie gewasse, soos aangedui. Ook vir die beheer van verderflike dopluis, op rustende rose.

If medical advice is needed, have product container or label at hand.

Active ingredients / Aktiewe bestanddele:

Sulphur.....800 g/kg

Registered by:

BASF South Africa (Pty) Ltd
Co. Reg. No. 66/10235/07
852 Sixteenth Road
P.O. Box 2801
Halfway House 1685
+27 11 203 2400

Geregistreeer deur:

BASF Suid-Afrika (Edms) Bpk
Mpy Reg. Nr 66/10235/07
852 Sestiende Laan
Posbus 2801
Halfweghuis 1685
+27 11 203 2400

Batch No. & Date manufactured: Refer to details printed on container
Lotnommer & Datum vervaardig: Verwys na besonderhede gedruk op houer

EMERGENCY NUMBER / NOODNOMMER: +27 82 446 8946



25 kg

® = Registered trademark of BASF / Geregistreeerde handelsmerk van BASF



81164036 ZA 2092



4 014348 730327