

<b>WARNINGS:</b>
<b>Hazard Statements</b>
<ul style="list-style-type: none"><li>Harmful if swallowed.</li> <li>Very toxic to aquatic life with long lasting effects.</li></ul>

- MAIZE:** ALLOW **90** DAYS BETWEEN PLANTING AND FEEDING.
- SUNFLOWER:** ALLOW **28** DAYS BETWEEN PLANTING AND HARVEST.

- Handle with care.
- Poisonous when absorbed through the skin or swallowed.
- Toxic to wildlife.
- Highly toxic to bees and other pollinators.
- TREATED SEED MAY BE POISONOUS TO SEED-EATING BIRDS. TREATED SEED MUST BE PLANTED PROPERLY AND TOTALLY COVERED WITH SOIL. NO SEED SHOULD BE LEFT IN PLACES WHERE IT CAN BE EATEN BY BIRDS.

- Sufficient soil moisture is necessary for the uptake of the active ingredient during germination.
- Use **Poncho® Votivo®** only for the treatment of high-quality certified seed.
- Do not use **Poncho® Votivo®** on sweet corn.
- Do not treat seed of inbred parent plants of hybrids, experimental hybrids or cultivars or newly released hybrids or cultivars with **Poncho® Votivo®**, without first referring to the manufacturer or seed suppliers.

- Planter should be calibrated prior to planting with treated seed to ensure that the required plant population is obtained. Calibration should be checked at regular intervals thereafter.
- Emergence of seedlings may be adversely affected if planting depth exceeds 50 mm. Efficacy on maize may also be adversely affected by planting depth.
- Treat only seed that is to be used for planting purposes. Never use treated seed for human or animal consumption.
- Store in original container in a cool, dry and well-ventilated place away from heat and sources of ignition and food and feedstuffs.
- Store in a place accessible by authorized persons only.
- Keep out of reach of children, uninformed persons and animals.
- In case of poisoning CALL A DOCTOR AND MAKE LABEL AVAILABLE TO HIM/HER.
- ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL SOIL, CLIMATIC AND STORAGE CONDITIONS AND/OR EXCESSIVELY HIGH INSECT/ PEST POPULATION PRESSURE; COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE PEST AGAINST THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

**PRECAUTIONS:**

**Precautionary Statements:**

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read label before use.
- Wear protective gloves/clothing/eye protection.
- Do not eat, drink or smoke when using this product.
- Wash contaminated body parts thoroughly after handling.
- IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.
- Rinse mouth
- Collect spillage.
- Dispose of contents/container to hazardous or special waste collection point.

- Avoid contact with spilled product or contaminated surfaces.
- Remove all sources of ignition.
- Do not allow product to get into surface water, drains and ground water.
- Destroy empty container by perforation and flattening and do not use for any other purpose.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.
- Handle treated seed carefully and clearly mark the bags which contain treated seed "POISONOUS" and never use this bag for any other purpose.

Hazardous ingredients	CAS number	Concentration (W/W)
According to UN GHS criteria		
clothianidin (ISO); 3-[(2-chloro-1,3-thiazol-5-yl)methyl]-2-methyl-1-nitroguanidine	210880-92-5	39,7 <span> </span> %
Bacillus firmus I-1582		8 <span> </span> %
1,2-Benzisothiazol-3(2H)-one	2634-33-5	< 0,05 <span> </span> %
mixture of: 5-chloro-2-methyl-2H-isothiazol-3-one and 2-methyl-2H-isothiazol-3-one (3:1)	55965-84-9	< 0,0015 <span> </span> %
Sodium chloride	7647-14-5	< 5 <span> </span> %
Glycerol	56-81-5	< 10 <span> </span> %
Propane-1,2-diol	57-55-6	< 5 <span> </span> %

**FIRST AID TREATMENT**

Remove contaminated clothing. If symptoms persist, seek medical advice. Move person to fresh air.

<b>ON CONTACT WITH EYES:</b>	Wash affected eyes for at least 15 minutes under running water with eyelids held open. Remove contact lenses, if present. If symptoms persist, seek medical advice.
<b>ON INGESTION:</b>	Rinse mouth immediately with water. Do not induce vomiting. Seek medical attention.
<b>ON SKIN CONTACT:</b>	Wash thoroughly with soap and water.
<b>IF INHALED:</b>	Keep patient calm, remove to fresh air, seek medical attention.

**TREATMENT:** Treat according to symptoms (decontamination, vital functions), no known specific antidote.

**RESISTANCE MANAGEMENT:**

For resistance management, **Poncho® Votivo®** is a group code 4A insecticide. Any insect population may contain individuals naturally resistant to **Poncho® Votivo®** and other group code 4A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **Poncho® Votivo®** or any other group code 4A insecticides.

To delay resistance:

- Avoid exclusive repeated use of insecticides from the same insecticide group code. Alternate with products from different insecticide group codes.
- Integrate other control methods (chemical, cultural, biological) into insect control programs.

**FOR SPECIFIC INFORMATION ON RESISTANCE MANAGEMENT, CONTACT THE REGISTRATION HOLDER, BASF SOUTH AFRICA (PTY) LTD.**

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED

**Storage Instructions:**

For best results **Poncho® Votivo®** must be stored between 0 °C - 30 °C. Ideally long-term product storage should have temperature-controlled conditions.

**Mixing Instructions:**

Mix the required volume of **Poncho® Votivo®** thoroughly with the required volume of water to make up a slurry volume equivalent to 1000 ml/100 kg seed before treating the seed.

All slurry mixtures that contain **Poncho® Votivo®** should be used within 72 hours of mixing. Large quantities of seed are conveniently treated in automatic seed dressing machines. The machines should be checked at intervals to ensure that the correct quantity of **Poncho® Votivo®** is used.

Maize seed may be treated with **Poncho® Votivo®** immediately before planting but can be treated earlier and stored under cool dry conditions for a season without germination being affected.

**APPLICATION TABLE:**

Crop	Dosage	Notes
<b>Sunflower</b>	0.96 ml/ 1000 seed	For soil insects as listed under <b>SOIL INSECTS CONTROLLED.</b> Treat seed just prior to planting.
<b>Maize</b>	0.96 ml/ 1000 seed	For soil insects as listed under <b>SOIL INSECTS CONTROLLED.</b> <p><b>Bio-stimulation:</b> Improved plant growth and root development, enhancing tolerance against stress and reducing early nematode* attack resulting in yield increase under certain conditions.</p> <ul style="list-style-type: none"><li>All above PLUS Leafhopper (<i>Cicadulina mbila</i>), the vector of streak disease.</li></ul> <p><b>Peridiam® EV26001</b> (Reg. No. L8669) at the registered rate (please refer to the <b>Peridiam® EV26001</b> product label) <b>MUST</b> be added to the <b>Poncho® Votivo®</b>.</p>
	1.68 ml/ 1000 seed	

**SOIL INSECTS CONTROLLED:**

Black maize beetles (Heteronychus arator),

Astylus larvae (Astylus atromaculatus),

Ground weevils (Protostrophus spp.),

Wireworms (Elateridae),

Lesser false wireworms/Dusty surface beetles (Gonocephalum and mesomorphus spp.),

Greater false wireworms (Somaticus spp.)

**NOTE:**

**Black maize beetle and ground weevil:**

Notwithstanding the use of this product, some black maize beetle and ground weevil damage could still be expected. Under conditions of very high infestation pressure 20 % or more of the maize crop may be damaged or destroyed, in spite of having been treated. More insect damage can be expected in soils with high levels of plant material associated with minimum tillage practices.

\*It is important to note that additional nematode control measures are advisable under conditions of high nematode pressure.

**Leafhopper:**

**Poncho® Votivo®** controls the leafhopper, which is the vector of the maize streak disease virus (MSV). Notwithstanding the use of this product some MSV damage could still be expected and therefore it is important to note that additional precautions must be taken in areas with high leafhopper populations.

The additional precautions includes:

- Consult your seed representative regarding a tolerant cultivar recommendation.
- Avoid exceptionally early or late plantings.
- Ensure good grass weed control in and around the maize field.
- Alternative precautions must also be taken in areas with high leafhopper population pressure, not only at the time of the maize planting but also on other crops in the area harboring leafhoppers.

**Risk management is based on a combination of the following measures:**

- Management practices:
  - Planting date and crop rotation can reduce the incidence of MSV.
  - Weed control to reduce the vector source.
  - Increased fertilizer application can also reduce MSV losses.
- Resistance is particularly effective and more visible in hybrid cultivars bred and developed in areas where the disease is endemic.
- Chemical control.

**CONDITIONS OF SALE AND WARRANTY**

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of this product in a manner inconsistent with its labeling, all of which are beyond the control of BASF South Africa (Pty) Ltd. ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**
BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of the Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

**Poncho® Votivo®** are registered trademarks of BASF.

**Peridiam®** is a registered trademark of the Bayer Group.

READ ATTACHED PACKAGE OR LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS



# Poncho® Votivo®

Reg. No. L9250 (Act No. 36 of 1947)

Reg. Nr. L9250 (Wet Nr. 36 van 1947)

A systemic insecticidal and biological seed dressing for use on maize and sunflower seed for the control of soil insects as specified.

'n Sistemiese insekdoder en biologiese saadbehandelingsmiddel vir gebruik op mielie- en sonneblomsaad vir die beheer van grond insekteplae soos aangedui.

Harmful if swallowed.

Very toxic to aquatic life with long lasting effects.

Do not eat, drink or smoke when using this product.

Wash contaminated body parts

thoroughly after handling.

GROUP	4A	INSECTICIDE
-------	----	-------------

<b>Active ingredients:</b>		<b>Aktiewe bestanddele:</b>
Clothianidin Neonicotinoid (chloro-nicotinyl)	500 g/l	Clothianidin Neonicotinoid (chloro-nicotinyl)
<i>Bacillus firmus</i>	100 g/l	<i>Bacillus firmus</i>

BATCH NO: / DATE OF MANUFACTURE: Refer details printed on container

LOTNOMMER: / DATUM VERVAARDIG: Verwys na besonderhede gedruk op houer

NET CONTENTS:	200 ℓ
NETTO INHOUD:	

DATE OF MANUFACTURE:

DATUM VERVAARDIG:

Registered by:	Geregistreer deur:
BASF South Africa (Pty) Ltd	BASF Suid-Afrika (Edms.) Bpk.
Reg. No. 1966/10235/07	Reg. Nr. 1966/10235/07
P.O. Box 2801	Posbus 2801
Halfway House, 1685	Halfway House, 1685
SOUTH AFRICA	SUID-AFRIKA
+27 11 203 2400	+27 11 203 2400

UN Number:	3082	VN Nommer
EMERGENCY NUMBER:	+27 82 446 8946	NOODNOMMER

**Poncho® Votivo®** are registered trademarks of BASF / is geregistreerde handelsmerke van BASF

# 200 ℓ

81162663 NVA 2023-05-0603-0225

**WAARSKUWINGS:**

<b>Gevaarstellings:</b>
<ul style="list-style-type: none"><li>Skadelik indien ingesluk.</li> <li>Baie giftig vir akwatiese organismes met langdurige effek.</li></ul>

- MIELIES:** LAAT **90** DAE TOE TUSSEN PLANTTYD EN VOER.
- SONNEBLOM:** LAAT **28** DAE TOE TUSSEN PLANTTYD EN VOER.
- Hanteer versigtig.
- Giftig wanneer ingeneem of deur die vel geabsorbeer.
- Giftig vir wild lewe.
- Utters giftig vir heuning bye en ander bestuivers.
- BEHANDELDE SAAD KAN GIFTIG WEES VIR SAADVRETNENDE VOËLS. BEHANDELDE SAAD MOET BEHOORLIK GEPLANT WORD EN HEELTEMAL MET GROND BEDEK WORD. GEEN SAAD MOET OP PLEKKE GELAAT WORD WAAR DIT DEUR VOËLS GEVREET KAN WORD NIE.

- Voeldoende grondvog is nodig vir die opname van die aktiewe bestanddeel gedurende ontkieming.
- Gebruik **Poncho® Votivo®** slegs vir behandeling van hoë gehalte gesertifiseerde saad.
- Moet nie **Poncho® Votivo®** op suiker mielies gebruik nie.
- Moet nie **Poncho® Votivo®** op saad van ingeteelde ouerplante van basters, eksperimentele basters of cultivars of nuut vrygestelde basters of cultivars gebruik, voordat die vervaardiger of saad voorsieners geraadpleeg is nie.
- Die planter moet voor plant in aanvang neem gekalibreer word met behandelde saad om te verseker dat die benodigde plantpopulasie verkry sal word. Die kalibrasie moet daarna met gereelde intervalle nagegaan word.
- Die opkoms van saalinge mag nadelig beïnvloed word indien saad dieper as 50 mm geplant word. Effektiviteit mag ook nadelig deur plantdiepte beïnvloed word op mielies.
- Behandel slegs saad wat geplant gaan word. Moet nooit behandelde saad as voedsel of voer gebruik nie.
- Bewaar in oorspronklike houers in 'n koel, droë en goed geventileerde plek weg van hitte en bronne wat ontbranding kan veroorsaak en voedsel en voer.
- Stoor in 'n plek wat slegs deur gemagtigde persone toeganklik is.
- Hou buite bereik van kinders, oningeligte persone en diere.
- In geval van vergiftiging ONTBID 'N GENEESHEER EN TOON ETIKET AAN HOM/HAAR. ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GETOETS IS WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE AANGESIEN DIE WERKING EN EFFEK DAARVAN BEÏNVLOED KAN WORD DEUR FAKTORE SOOS ABNORMALE GROND-, KLIMAATS- EN BERGINGSTOESTANDE EN/OF UITERMATIGE HOË INSEK/PES POPULASIE DRUK; VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE EN DIE VOORKOMS VAN WEERSTAND VAN DIE PLAAG TEEN DIE BETROKKE MIDDEL SOWEL AS DIE METODEDE, TYD EN AKKURAATHEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADELIGE EFFEK OP MENS OF DIER OF VIR 'N GEBREK AAN PRESTASIE AS GEVOLG VAN DIE VERSUIM VAN DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID. VOORSORGMAAATREËLS:

**Voorsorgstellings:**

- Indien mediese advies benodig word, hou die produkhouer of -etiket byderhand.
- Hou buite bereik van kinders.
- Lees sorgvuldig en volg alle instruksies.
- Dra beskermende handskoene en oogbeskerming of gesigbeskerming.
- Moenie eet rook of drink terwyl produk hanteer word nie.
- Was gekontamineerde liggaamsdele deeglik na hantering.
- INDIEN INGESLUK: Skakel 'n GIFBEHEERSENTRUM of dokter/genesheer indien u siek voel.
- Spoel mond uit.
- Maak stortslu bymekaar.
- Raak ontslae van inhoud en houer deur na 'n gevaarlike- of spesiale-afval insamelsplek te besorg.

- Vermy kontak met verspilde produk of gekontamineerde oppervlakte.
- Verwyder alle bronne wat ontbranding kan veroorsaak.
- Moet nie toelaat dat produk in oppervlakte water, dreineer stelsels en grond water kom nie.
- Spoel leë houer drie keer uit met 'n volume water gelyk aan ten minste een tiende van die van die houer en voeg die uitspoelwater by die inhoud van die spuittenk voordat van die houer op die voorgeskrewe metode ontslae geraak word. Dra sorg dat die volume water wat benodig word vir behandeling nie oorskry word nie.
- Vernietig leë houer deur gate daarin te maak en plat te slaan. Moet vir geen doel weer gebruik nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- Hanteer behandelde saad versigtig. Merk sakke wat behandelde saad bevat duidelik "GIFTIG" en moet hulle nooit vir enige ander doel gebruik nie.

**RELEVANTE STOWWE**

Gevaarlike bestanddele (Volgens UN GHS kriteria)	CAS nommer	Konsentrasie (W/W)
clothianidin (ISO); 3-[(2-chloro-1,3-thiazol-5-yl)methyl]-2-methyl-1-nitroguanidine	210880-92-5	39,7 <span> </span> %
Bacillus firmus I-1582		8 <span> </span> %
1,2-Benzisothiazol-3(2H)-one	2634-33-5	< 0,05 <span> </span> %
mixture of: 5-chloro-2-methyl-2H-isothiazol-3-one and 2-methyl-2H-isothiazol-3-one (3:1)	55965-84-9	< 0,0015 <span> </span> %
Sodium chloride	7647-14-5	< 5 <span> </span> %
Glycerol	56-81-5	< 10 <span> </span> %
Propane-1,2-diol	57-55-6	< 5 <span> </span> %

**NOODHULPEBEHANDELING**

Verwyder besoedelde klere. Indien simptome voortduur, verkry mediese hulp. Verwyder persoon na vars lug

<b>INDIEN IN DIE OË:</b>	Spoel aangetaste oë vir ten minste 15 minute met lopende water, terwyl die ooglede oopgehou word.
<b>INDIEN INGESLUK:</b>	Spoel mond dadelik uit met water. Moenie braking veroorsaak nie. Verkry mediese hulp
<b>INDIEN OP DIE VEL:</b>	Was vel deeglik met seep en water.

**INDIEN INGEASEM:** Hou pasiënt kalm en verwyder na vars lug. Verkry mediese hulp.

**BEHANDELING:** Behandel simptomaties (reinig, belangrikste funksies). Daar is geen spesifieke teenmiddel nie.

**WEERSTANDBESTUUR:**

**Poncho® Votivo®** is 'n groepkode 4A insekdoder. Enige populasie van 'n spesifieke insek mag individue insluit wat 'n natuurlike weerstand teen **Poncho® Votivo®** of enige ander groepkode 4A insekdoder het. Indien hierdie insekdoders herhaaldelik aangewend word kan die weerstandbiedende individue uiteindelik die insekpopulasie oorheers. Hierdie weerstandbiedende insekte sal waarskynlik nie deur **Poncho® Votivo®** of enige ander groepkode 4A insekdoder beheer word nie.

Om weerstand teen insekdoders te verraag:

- Vermy die eksklusiewe herhaaldelike gebruik van insekdoders in dieselfde insekdoder groepkode.
- Wissel af met produkte in verskillende insekdoder groepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in insekbeheer programme.

Vir spesifieke inligting oor weerstandbestuur kontak die registrasiehouer van hierdie produk.
**GEBRUIKSAANWYSINGS:** GEBRUIK SLEGS SOOS AANBEVEEL.

**Bergingsinstruksies:**

Vir die beste resultate behoort **Poncho® Votivo®** teen 0 °C - 30 °C opgeberg te word. Lang termyn opberging van die produk behoort onder temperatuur gereguleerde toestande te geskiet.

**Menginstruksies**

Meng die benodigde volume **Poncho® Votivo®** deeglik met die benodigde volume water om 'n totale mengsel volume van 1000 ml /100 kg saad op te maak voordat die saad daarmee behandel word.

Alle opgemaakte “slurries” (**Poncho® Votivo®** + water) moet binne 72 uur na opmaak gebruik word.

Groot hoeveelhede saad kan doeltreffend in outomatiese saadbehandelingsmasjiene behandel word. Die masjiën moet gereeld nagegaan word om te verseker dat die regte hoeveelheid **Poncho® Votivo®** toegedien word.

Mieliesaad kan onmiddellik voor planting met **Poncho® Votivo®** behandel word of vroeër en dan vir 'n seisoen lank opgeberg word onder koel droë toestande sonder dat ontkieming benadeel sal word.

**TOEDIENINGSTABEL:**

Gewas	Dosis	Notas
<b>Sonneblom</b>	0.96 ml/ 1000 pitte	Vir grondinsekte soos gelys onder <b>GRONDINSEKTE WAT BEHEER WORD</b> <p>Behandel saad net voor plant</p>
<b>Mielies</b>	0.96 ml/ 1000 pitte	Vir grondinsekte soos gelys onder <b>GRONDINSEKTE WAT BEHEER WORD</b> <p><b>Bio-stimulasie:</b> Verbeterde plant groei en wortel ontwikkeling wat die verdraagsaamheid teenoor stres verbeter en vroeë aalwurm aanvalle verminder met gevolglike verhoogde oes opbrengs onder sekere omstandighede. Egogenoemde PLUS Blaarspringer (<i>Cicadulina mbila</i>), die vektor van streepsiekte</p> <p><b>Peridiam® EV26001</b> (Reg. Nr. L8669) teen die geregistreerde dosis (verwys asseblief na die <b>Peridiam® EV26001</b> produk etiket) <b>MOET</b> by die <b>Poncho® Votivo®</b> gevoeg word.</p>

**GRONDINSEKTE WAT BEHEER WORD**

- Astylus larwes (*Astylus atromaculatus*)
- Swartmielekewers (*Heteronychus arator*),
- Kleinvaldraadwurms /Vaalgrondkewers (*Gonocephalum and Mesomorphus spp.*),
- Draadwurms (*Elateridae*)
- Grootvaldraadwurms (*Somaticus spp.*)
- Grondkalanders (*Protostrophus spp.*)

**NOTA:**

**Swartmielekewer en grondkalander:**

Nieteenstaande die gebruik van hierdie middel, kan swartmielekewer en grondkalander beskadiging nog steeds verweg word. Onder toestande van hoë infestasiedruk kan selfs 20 % of meer van die mielies ten spyte van behandeling, beskadig of vernietig word. Meer insekskade kan verweg word in grade met hoë vlakke van plantmateriaal geassosieer met minimum bewerkings praktyke.

\* Dit is belangrik om daarop te let dat bykomende aalwurmbeheermaatreëls raadsaam is onder toestande van hoë aalwurm druk.

**Blaarspringer:**

**Poncho® Votivo®** beheer die blaarspringer wat die vektor van die mieliestreepsiekte virus (MSV) is. Nieteenstaande die gebruik van hierdie middel kan MSV beskadiging nog steeds verweg word en daarom is dit belangrik om daarop te let dat bykomende voorsorgmaatreëls getref moet word in areas met hoë blaarspringer populasies.

Die voorsorgmaatreëls sluit in:

- Raadpleeg u saadverteenwoordiger rakende 'n verdraagsame kultivar aanbeveling.
- Vermy buitengewoon vroeë of laat aanplantings.
- Verseker goeie gras onkruidbeheer in en om die mielieland.
- Alternatiewe voorsorgmaatreëls moet ook getref word in areas met hoë blaarspringer populasie druk